

# Ergodicity in statistical mechanics of interacting and disordered systems: Destroying and restoring equilibrium ergodic states

Václav Janiš

Institute of Physics, Academy of Sciences of the Czech Republic, Praha, CZ  
and Fulbright Scholar at Louisiana State University, Baton Rouge, USA

20th Mardi Gras Conference,  
LSU, February 15, 2015

Collaborators: Anna Kauch, Antonín Klíč (FZÚ AV ČR, v. v. i.)

# Outline

- 1 Introduction - ergodicity and thermodynamic homogeneity
- 2 Ergodicity breaking in spin models
  - Models of interacting spins
  - Models with disorder and frustration - spin glasses
- 3 Real-replica method for disordered frustrated systems
  - Replication of the phase space
  - Discrete replica-symmetry (replica-independence) breaking
  - Continuous replica-symmetry breaking
- 4 Application: Solvable cases
  - Replica symmetric and one-level RSB -- Ising
  - Infinite RSB - asymptotic solution -- Ising
  - Potts and  $p$ -spin glass
- 5 Conclusions

# Microscopic dynamics of large systems

## Instantaneous microscopic dynamics of large systems

- Macroscopic objects -- aggregate of microscopic elements
- Superposition principle -- large system of coupled Hamilton equations determining the phase-space trajectory
- **Impossibility to determine complete initial conditions**
- Inability to determine the actual trajectory of large microscopic states
- **Macroscopic state:** phase space fluid with Liouville equation
- Entropy as a measure of macroscopic uncertainty

How do we determine macroscopic properties without solving Liouville equation?

# Macroscopic time scales

## Large time scales - macroscopic stationarity

- Macroscopic measurements -- on large time scales (relaxation time)
- Time fluctuations on microscopic time scales macroscopically unimportant
- **Only time averaged quantities measurable** (relevant)
- Energy as the only relevant macroscopically conserved quantity restriction on the phase-space trajectory
- **Thermodynamic equilibrium** -- macroscopically static state

How do we calculate time averaged quantities?

# Ergodicity in equilibrium statistical physics

- **Fundamental ergodic theorem** (Birkhoff)

$$\langle f \rangle_T \equiv \lim_{T \rightarrow \infty} \frac{1}{T} \int_{t_0}^{t_0+T} f(X(t)) dt = \frac{1}{\Sigma_E} \int_{S_E} f(X) dS_E \equiv \langle f \rangle_S$$

- Phase space homogeneously covered by the phase trajectory

$$\lim_{T \rightarrow \infty} \frac{\tau_R}{T} = \frac{\Sigma_R(E)}{\Sigma(E)}$$

- **Equilibrium ergodic macroscopic state**
  - homogeneously spread over the allowed phase space
  - characterized by homogeneous parameters ( $\{E, T\}, \{N, \mu\}, \dots$ )
  - **number of relevant parameters (Legendre pairs) a priori unknown**

How do we determine the phase space covered by the phase space trajectory?

# Homogeneity of thermodynamic potentials

- Homogeneity in the phase space

$$S(E) = k_B \ln \Gamma(E) = \frac{k_B}{\nu} \ln \Gamma(E)^\nu = \frac{k_B}{\nu} \ln \Gamma(\nu E)$$

$$F(T) = - \frac{k_B T}{\nu} \ln [\text{Tr} e^{-\beta H}]^\nu = - \frac{k_B T}{\nu} \ln [\text{Tr} e^{-\beta \nu H}]$$

- Homogeneity of thermodynamic potentials (Euler)

$$\alpha F(T, V, N, \dots, X_i, \dots) = F(T, \alpha V, \alpha N, \dots, \alpha X_i, \dots)$$

Density of the free energy  $f = F/N$

-- function of only **densities** of extensive variables  $X_i/N$

Ergodicity (homogeneity) guarantees existence and uniqueness of the thermodynamic limit  $N \rightarrow \infty$

# Homogeneity of thermodynamic potentials

- Homogeneity in the phase space

$$S(E) = k_B \ln \Gamma(E) = \frac{k_B}{\nu} \ln \Gamma(E)^\nu = \frac{k_B}{\nu} \ln \Gamma(\nu E)$$

$$F(T) = - \frac{k_B T}{\nu} \ln [\text{Tr} e^{-\beta H}]^\nu = - \frac{k_B T}{\nu} \ln [\text{Tr} e^{-\beta \nu H}]$$

- Homogeneity of thermodynamic potentials (Euler)

$$\alpha F(T, V, N, \dots, X_i, \dots) = F(T, \alpha V, \alpha N, \dots, \alpha X_i, \dots)$$

Density of the free energy  $f = F/N$

-- function of only **densities** of extensive variables  $X_i/N$

Ergodicity (homogeneity) guarantees existence and uniqueness of the thermodynamic limit  $N \rightarrow \infty$

# Ergodicity breaking

- Ergodicity gives meaning to statistical averages
- Thermodynamic properties in the infinite-volume limit
- Ergodicity breaking -- improper statistical phase space
  - 1 caused by a phase transition breaking a symmetry of the Hamiltonian
  - 2 without apparent symmetry breaking -- glass-like behavior
- Means to restore ergodicity
  - 1 Measurable (physical) symmetry breaking fields
  - 2 Real replicas (non-measurable symmetry breaking fields)

Ergodicity must be restored to establish stable equilibrium

# Ergodicity breaking

- Ergodicity gives meaning to statistical averages
- Thermodynamic properties in the infinite-volume limit
- Ergodicity breaking -- improper statistical phase space
  - 1 caused by a phase transition breaking a symmetry of the Hamiltonian
  - 2 without apparent symmetry breaking -- glass-like behavior
- Means to restore ergodicity
  - 1 Measurable (physical) symmetry breaking fields
  - 2 Real replicas (non-measurable symmetry breaking fields)

Ergodicity must be restored to establish stable equilibrium

- Model of interacting spins (Heisenberg)

$$H[J, \mathbf{S}] = - \sum_{i < j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j$$

- Spin exchange

- Ferromagnetic interaction:  $J_{ij} > 0$
- Antiferromagnetic interaction:  $J_{ij} < 0$

- Regular crystalline structure (lattice)

- Strong anisotropy: Only single spin projection ( $S^z$ ) interacts

- Ising model

$$H[J, S] = - \sum_{i < j} J_{ij} S_i S_j$$

- Classical spins with  $S_i = \pm 1$  ( $\hbar/2$  units)

# Other spin models -- generalizations of Ising I

- Potts model -  $p > 2$  spin projections


$$H_p = - \sum_{i < j} J_{ij} \delta_{n_i, n_j}$$

$$n_i = 1, 2, \dots, p$$

- Spin representation

$$H_p [J, \mathbf{S}] = -\frac{1}{2} \sum_{i,j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j - \sum_i \mathbf{h} \cdot \mathbf{S}_i,$$

Potts vectors  $\mathbf{S}_i = \{s_i^1, \dots, s_i^{p-1}\}$ , values are state vectors  $\{\mathbf{e}_A\}_{A=1}^p$



# Other spin models -- generalizations of Ising II

$$\sum_{A=1}^p e_A^\alpha = 0, \quad \sum_{A=1}^p e_A^\alpha e_A^\beta = p \delta^{\alpha\beta}, \quad e_A^\alpha e_B^\alpha = p \delta_{AB} - 1$$

## ■ Explicit representation

$$e_A^\alpha = \begin{cases} 0 & A < \alpha \\ \sqrt{\frac{p(p-\alpha)}{p+1-\alpha}} & A = \alpha \\ \frac{1}{\alpha-p} \sqrt{\frac{p(p-\alpha)}{p+1-\alpha}} & A > \alpha. \end{cases}$$

## ■ $p$ -spin model

$$H_p[J, S] = \sum_{1 \leq i_1 < i_2 < \dots < i_p} J_{i_1 i_2 \dots i_p} S_{i_1} S_{i_2} \dots S_{i_p}.$$

$S$  are Ising spins,  $p = 2$  reduces to Ising

# Ising thermodynamics -- mean-field solution

- Thermally induced spin fluctuations -- free energy

$$-\beta F(T) = \ln \text{Tr}_S \exp \{-\beta H[J, S]\}$$

- Long-range ferromagnetic interaction:  $J_{ij} = -J/N$
- Mean-field (Weiss) solution

$$F(T, m)/N = \frac{Jm^2}{2} - \frac{1}{\beta} \ln 2 \cosh(\beta Jm)$$

with a global magnetization  $m$



- Equilibrium state -- magnetization minimizing  $F(T, m)$
- Equilibrium magnetization

$$m = \tanh(\beta Jm)$$



# Disorder & frustration -- inhomogeneous spin exchange

- Randomness in the spin exchange
- **System locally frustrated**: ferro (red bond) and antiferro (blue bond) randomly distributed





# Disorder & frustration -- inhomogeneous spin exchange

## III

- Potts model (not symmetric v.r.t. spin reflection)

$$P(J_{ij}) = \sqrt{\frac{N}{2\pi J^2}} \exp \frac{-N(J_{ij} - J_0/N)^2}{2J^2},$$

$J_0 = \sum_j J_{0j}$  -- averaged (ferromagnetic) interaction

- $p$ -spin model

$$P(J_{i_1 i_2 \dots i_p}) = \sqrt{\frac{N^{p-1}}{\pi p!}} \exp \left\{ -\frac{J_{i_1 i_2 \dots i_p}^2 N^{p-1}}{J^2 p!} \right\}$$





# Real replicas -- stability w.r.t. phase-space scalings

Real replicas -- means to probe thermodynamic homogeneity

Replicated Hamiltonian:  $[H]_\nu = \sum_{a=1}^{\nu} H^a = \sum_{\alpha=1}^{\nu} \sum_{\langle ij \rangle} J_{ij} S_i^{\alpha} S_j^{\alpha}$

Symmetry-breaking fields:  $\Delta H(\mu) = \frac{1}{2} \sum_{a \neq b} \sum_i \mu^{ab} S_i^a S_i^b$

Averaged replicated free energy with coupled replicas

$$F_\nu(\mu) = -k_B T \frac{1}{\nu} \left\langle \ln \text{Tr} \exp \left\{ -\beta \sum_a^{\nu} H^a - \beta \Delta H(\mu) \right\} \right\rangle_{av}$$

Analytic continuation to non-integer parameter  $\nu$

Stability w.r.t. phase space scaling:

$$\lim_{\mu \rightarrow 0} \frac{dF_\nu(\mu)}{d\nu} \equiv 0$$

Real replicas - simulate impact of surrounding bath

# Real replicas -- stability w.r.t. phase-space scalings

Real replicas -- means to probe thermodynamic homogeneity

Replicated Hamiltonian:  $[H]_\nu = \sum_{a=1}^{\nu} H^a = \sum_{\alpha=1}^{\nu} \sum_{\langle ij \rangle} J_{ij} S_i^{\alpha} S_j^{\alpha}$

Symmetry-breaking fields:  $\Delta H(\mu) = \frac{1}{2} \sum_{a \neq b} \sum_i \mu^{ab} S_i^a S_i^b$

Averaged replicated free energy with coupled replicas

$$F_\nu(\mu) = -k_B T \frac{1}{\nu} \left\langle \ln \text{Tr} \exp \left\{ -\beta \sum_a^{\nu} H^a - \beta \Delta H(\mu) \right\} \right\rangle_{av}$$

Analytic continuation to non-integer parameter  $\nu$

Stability w.r.t. phase space scaling:

$$\lim_{\mu \rightarrow 0} \frac{dF_\nu(\mu)}{d\nu} \equiv 0$$

Real replicas - simulate impact of surrounding bath

# Annealed vs. quenched disorder

- Averaged ( $\nu$ -times replicated) partition function

$$\langle Z_N^\nu \rangle_{av} = \int D[J] \mu[J] \prod_{a=1}^{\nu} \prod_{i=1}^N d[\mathbf{S}_i^a] \rho[\mathbf{S}_i^a] \exp \left\{ -\beta \sum_{a=1}^{\nu} H[J, \mathbf{S}^a] \right\}$$

- Averaged ( $\nu$ -times replicated) free energy

$$-\beta \langle F_N^\nu \rangle_{av} = \int D[J] \mu[J] \ln \int \prod_{a=1}^{\nu} \prod_{i=1}^N d[\mathbf{S}_i^a] \rho[\mathbf{S}_i^a] \exp \left\{ -\beta \sum_{a=1}^{\nu} H[J, \mathbf{S}^a] \right\}$$

- Replicas for disordered systems:

- **Quenched disorder** (spin glasses) -- replica trick ( $\nu \rightarrow 0$ )

$$\beta F_{qu} = - \lim_{\nu \rightarrow 0} \left[ \frac{1}{\nu} \lim_{N \rightarrow \infty} (\langle Z_N^\nu \rangle_{av} - 1) \right]$$

- **Annealed disorder** -- thermodynamic homogeneity ( $\nu$  arbitrary)

$$\beta F_{an} = - \frac{1}{\nu} \lim_{N \rightarrow \infty} \ln \langle Z_N^\nu \rangle_{av}$$

Copyright © 2015, Laurent Micco, all rights reserved. Copyright © 2015, Laurent Micco, all rights reserved.

# Ergodicity breaking -- broken LRT in replicated space

- Breaking of LRT to inter-replica interaction  $\mu^{ab} \rightarrow 0$

$$f_\nu = \frac{\beta \mathcal{J}^2}{4} \left[ \frac{1}{\nu} \sum_{a \neq b}^{\nu} \left\{ (\chi^{ab})^2 + 2q\chi^{ab} \right\} - (1-q)^2 \right]$$

$$- \frac{1}{\beta \nu} \int_{-\infty}^{\infty} \frac{d\eta}{\sqrt{2\pi}} e^{-\eta^2/2} \ln \text{Tr}_\nu \exp \left\{ \beta^2 \mathcal{J}^2 \sum_{a < b}^{\nu} \chi^{ab} S^a S^b + \beta \bar{h} \sum_{a=1}^{\nu} S^a \right\}$$

$$\chi^{ab} = \langle \langle S^a S^b \rangle_T \rangle_{av} - q, \quad q = \langle \langle S^a \rangle_T^2 \rangle_{av}, \quad \bar{h} = h + \eta \sqrt{q}$$

- Free energy  $f_\nu$  must be analytic function of index  $\nu$
- Parisi conditions for analytic continuation

$$\chi^{aa} = 0, \quad \chi^{ab} = \chi^{ba}, \quad \sum_{c=1}^{\nu} (\chi^{ac} - \chi^{bc}) = 0$$

- $K < \nu - 1$  different inter-replica susceptibilities  $\chi_1, \dots, \chi_K$  with multiplicities  $\nu_1, \dots, \nu_K$

# Ergodicity breaking -- broken LRT in replicated space

- Breaking of LRT to inter-replica interaction  $\mu^{ab} \rightarrow 0$

$$f_\nu = \frac{\beta \mathcal{J}^2}{4} \left[ \frac{1}{\nu} \sum_{a \neq b}^{\nu} \left\{ (\chi^{ab})^2 + 2q\chi^{ab} \right\} - (1-q)^2 \right]$$

$$- \frac{1}{\beta \nu} \int_{-\infty}^{\infty} \frac{d\eta}{\sqrt{2\pi}} e^{-\eta^2/2} \ln \text{Tr}_\nu \exp \left\{ \beta^2 \mathcal{J}^2 \sum_{a < b}^{\nu} \chi^{ab} S^a S^b + \beta \bar{h} \sum_{a=1}^{\nu} S^a \right\}$$

$$\chi^{ab} = \langle \langle S^a S^b \rangle_T \rangle_{av} - q, \quad q = \langle \langle S^a \rangle_T^2 \rangle_{av}, \quad \bar{h} = h + \eta \sqrt{q}$$

- Free energy  $f_\nu$  must be analytic function of index  $\nu$
- Parisi conditions for analytic continuation

$$\chi^{aa} = 0, \quad \chi^{ab} = \chi^{ba}, \quad \sum_{c=1}^{\nu} (\chi^{ac} - \chi^{bc}) = 0$$

- $K < \nu - 1$  different inter-replica susceptibilities  $\chi_1, \dots, \chi_K$   
with multiplicities  $\nu_1, \dots, \nu_K$

# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters –  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_2 & \dots & q_{\nu-2} & q_{\nu-1} & q_{\nu} \\ q_0 & 0 & q_1 & q_2 & \dots & q_{\nu-2} & q_{\nu-1} & q_{\nu} \\ q_1 & q_1 & 0 & q_0 & \dots & q_{\nu-2} & q_{\nu-1} & q_{\nu} \\ q_1 & q_1 & q_0 & 0 & \dots & q_{\nu-2} & q_{\nu-1} & q_{\nu} \\ \dots & \dots \\ q_{\nu-2} & q_{\nu-2} & q_{\nu-2} & q_{\nu-2} & \dots & 0 & q_0 & q_1 & q_2 \\ q_{\nu-1} & q_{\nu-1} & q_{\nu-1} & q_{\nu-1} & \dots & q_0 & 0 & q_1 & q_2 \\ q_{\nu} & q_{\nu} & q_{\nu} & q_{\nu} & \dots & q_1 & q_1 & 0 & q_0 \\ \dots & \dots & \dots & \dots & \dots & q_2 & q_2 & q_0 & 0 \end{pmatrix}$$

$$q_i = q + \chi_i, \quad \nu = 2^d, \quad \nu - 1 = \sum_{i=1}^d \nu_i$$

# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_i = q + \chi_i, \nu_i = 2^i, \nu - 1 = \sum_i^K \nu_i$$

# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_i = q + \chi_i, \nu_i = 2^i, \nu - 1 = \sum_i^K \nu_i$$

# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_l = q + \chi_l, \nu_l = 2^l, \nu - 1 = \sum_l^K \nu_l$$

# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_l = q + \chi_l, \nu_l = 2^l, \nu - 1 = \sum_l^K \nu_l$$

# Analytic continuation


- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_l = q + \chi_l, \nu_l = 2^l, \nu - 1 = \sum_l^K \nu_l$$


# ultrametric structure

- ultrametric structure
- only block matrices of identical elements
  - larger blocks multiples of smaller blocks
  - hierarchy of embeddings around diagonal
  - ultrametric metrics (tree-like)





# Multiple embeddings -- including boundary terms



- $\Delta\chi_l$  -- inter-replica interaction strength,  
 $\lambda_l$  -- effective magnetic field due to replicated spins
- $\nu_l V$ : volume affected by replicated spins -- range of inter-replica interaction

$$\frac{N}{V} \ln Z_{l-1}(\beta, \bar{h}_l)$$

$$\rightarrow \frac{N}{\nu_l V} \ln \int \mathcal{D}\lambda_l Z_{l-1}^{\nu_l}(\beta, \bar{h}_l + \lambda_l \sqrt{\Delta\chi_l})$$

- Effective weight of surrounding spins in thermal averaging

$$\rho_l = \frac{Z_{l-1}^{\nu_l}}{\langle Z_{l-1}^{\nu_l} \rangle_{\lambda_l}}$$

# One-level embedding (Ising)

- Local magnetization after thermalization of surrounding spins

$$m_i = \left\langle \rho^{(\nu)}(h + \eta_i; \lambda, \chi) \tanh[\beta(h + \eta_i + \lambda J \sqrt{\chi})] \right\rangle_{\lambda} \equiv \langle \rho_i^{\nu} t_i \rangle_{\lambda}$$

$$t \equiv \tanh \left[ \beta \left( h + \eta \sqrt{q} + \sum_{l=1}^K \lambda_l \sqrt{\Delta \chi_l} \right) \right] \text{ with}$$

$$\langle X(\lambda_l) \rangle_{\lambda_l} \equiv \int_{-\infty}^{\infty} \mathcal{D} \lambda_l X(\lambda_l)$$

- $\rho_i^{\nu} \equiv \rho^{(\nu)}(h + \eta_i; \lambda, \chi)$  -- spin density (in volume  $V_0$ ) including impact of the bath (volume  $V_1$ )
- Fluctuating internal magnetic field (Gaussian random)

$$\eta_i = \sum_j J_{ij} m_j - m_i \left[ \sum_j \beta J_{ij}^2 (1 - m_j^2) + \beta \mathcal{J}(\nu - 1) \chi \right]$$

- Red terms -- impact of thermalizing of outer spins (volume  $V_1$ )

# Equilibrium state -- stationarity equations & stability

- Stationarity equations with discrete  $K$  replica hierarchies

$$\begin{aligned}
 q &= \langle \langle t \rangle_K^2 \rangle_\eta, \\
 \Delta\chi_l &= \langle \langle \langle t \rangle_{l-1}^2 \rangle_K \rangle_\eta - \langle \langle \langle t \rangle_l^2 \rangle_K \rangle_\eta, \\
 \nu_l \Delta\chi_l &= \frac{4}{\beta^2} \frac{\langle \langle \ln Z_{l-1} \rangle_K \rangle_\eta - \langle \langle \ln Z_l \rangle_K \rangle_\eta}{2 \left( q + \sum_{i=l+1}^K \Delta\chi_i \right) + \Delta\chi_l}
 \end{aligned}$$

$$t \equiv \tanh \left[ \beta \left( h + \eta \sqrt{q} + \sum_{l=1}^K \lambda_l \sqrt{\Delta\chi_l} \right) \right],$$

$$\langle t \rangle_l(\eta; \lambda_K, \dots, \lambda_{l+1}) = \langle \rho_l \dots \langle \rho_1 t \rangle_{\lambda_1} \dots \rangle_{\lambda_l}, \quad \rho_l = Z_{l-1}^{\nu_l} / \langle Z_{l-1}^{\nu_l} \rangle_{\lambda_l}$$

- Stability conditions determine number  $K$

$$\Lambda_l^K = 1 - \beta^2 \left\langle \left\langle \left\langle 1 - t^2 + \sum_{i=0}^l \nu_i \left( \langle t \rangle_{i-1}^2 - \langle t \rangle_i^2 \right) \right\rangle_l \right\rangle_K \right\rangle_\eta \geq 0$$



# Infinite many replica hierarchies II

- Anti time-ordering product from quantum many-body PT

$$\bar{\mathbb{T}}_\lambda \exp \left\{ \int_0^1 d\lambda \hat{O}(\lambda) \right\} \equiv 1 + \sum_{n=1}^{\infty} \int_0^1 d\lambda_1 \int_0^{\lambda_1} \dots \int_0^{\lambda_{n-1}} d\lambda_n \hat{O}(\lambda_n) \dots \hat{O}(\lambda_1)$$

- Initial condition (1RSB)

$$g_1(h) \equiv g_1(m_1, h) = \frac{1}{m_1} \ln \int_{-\infty}^{\infty} \frac{d\phi}{\sqrt{2\pi}} e^{-\phi^2/2} [2 \cosh(\beta(h + \phi\sqrt{\chi_1}))]^{m_1}$$

- Closed implicit equation

$$\begin{aligned} g'_1(m, h) &= \mathbb{E}(m, m_1, h) \circ g_0(h) \\ &\equiv \bar{\mathbb{T}}_m \exp \left\{ \frac{1}{2} \int_m^{m_1} dm' x(m') [\partial_{\bar{h}}^2 + 2m' g'_1(m'; h + \bar{h}) \partial_{\bar{h}}] \right\} g'_1(h + \bar{h}) \Big|_{\bar{h}=0} \end{aligned}$$

- Notation:  $X_0(m) = \int_{m_0}^m dm' x(m')$

# Discrete vs. continuous replica-symmetry breaking

## Discrete RSB

- 1 Hierarchical embeddings -- ultrametric structure
- 2 No restriction on the replica-induced order parameters
- 3 The number of replica hierarchies  $K$  from stability conditions
- 4 Either unstable or locally stable

## Continuous RSB

- 1 Limit of infinite number of replica hierarchies
- 2 Infinitesimal distance between replica hierarchies
- 3 Closed theory independently of stability of the discrete scheme
- 4 Always marginally stable

# Replica symmetric solution -- ergodic assumption

- No replication, only original spins -- ergodic assumption
- Ergodicity not broken in the low-temperature phase with a single order parameter  $q = \langle m^2 \rangle$  (replica symmetric)
- Free energy ( $J^2 = 1$ )



$$f(q) = -\frac{\beta}{4}(1-q)^2 - \frac{1}{\beta} \int_{-\infty}^{\infty} \mathcal{D}\eta \ln 2 \cosh [\beta (h + \eta\sqrt{q})]$$

- Stationarity equation  $q = \int_{-\infty}^{\infty} \mathcal{D}\eta \tanh^2 [\beta (h + \eta\sqrt{q})]$
- Stability condition

$$\Lambda = 1 - \beta^2 \left\langle \left( 1 - \tanh^2 [\beta (h + \eta\sqrt{q})] \right)^2 \right\rangle_{\eta} \geq 0$$

- Zero-temperature entropy:  $S(0) = -\sqrt{\frac{2}{\pi}} \approx -0.798$

# Replica symmetric solution - thermodynamics



# First level replica-symmetry (ergodicity) breaking

- Ergodicity broken in the SG phase -- one embedding

$$f(q; \chi, \nu) = -\frac{\beta}{4}(1-q)^2 + \frac{\beta}{4}(\nu-1)\chi(2q+\chi) + \frac{\beta}{2}\chi \\ - \frac{1}{\beta\nu} \int_{-\infty}^{\infty} \mathcal{D}\eta \ln \int_{-\infty}^{\infty} \mathcal{D}\lambda \{2 \cosh[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})]\}^\nu$$

- Stationarity equations ( $t \equiv \tanh[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})]$ )



$$q = \langle \langle t^2 \rangle_\lambda \rangle_\eta, \quad q_{EA} = q + \chi = \langle \langle t^2 \rangle_\lambda \rangle_\eta \\ \beta^2 \chi(2q + \chi)\nu = [\langle \ln \cosh[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})] \rangle_\lambda \\ - \ln \langle \cosh^\nu[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})] \rangle_\lambda^{1/\nu}]$$

- Stability conditions

$$\Lambda_0 = 1 - \beta^2 \langle \langle (1-t)^2 \rangle_\lambda \rangle_\eta$$

$$\Lambda_1 = 1 - \beta^2 \langle \langle 1 - (1-\nu)t^2 - \nu \langle t^2 \rangle_\lambda \rangle_\lambda \rangle_\eta$$

# 1RSB - thermodynamics



google:Louis Potts, Rough, Lattice, Glass, pdf, google:Louis Potts, Rough, Lattice, Glass, pdf



# SK model at zero magnetic field

Only asymptotic expansions available for  $K \rightarrow \infty$

- Small expansion parameter  $\tau = (T_c - T)/T_c$

$$\Delta\chi_l^K \doteq \frac{2}{2K+1} \tau, \quad \nu_l^K \doteq \frac{4(K-l+1)}{2K+1} \tau, \quad q^K \doteq \frac{1}{2K+1} \tau,$$

$$Q^K \equiv q_{EA} = q + \chi_1 - \chi_K \doteq \tau + \frac{12K(K+1)+1}{3(2K+1)^2} \tau^2, \quad \Lambda_l^K \doteq -\frac{4}{3} \frac{\tau^2}{(2K+1)^2}$$

$$\chi_T \doteq \beta \left( 1 - Q^K + \sum_{l=1}^K m_l \Delta\chi_l \right) \doteq 1 - \frac{\tau^2}{3(2K+1)^2}$$

$$\Delta f \doteq \left( \frac{1}{6} \tau^3 + \frac{7}{24} \tau^4 + \frac{29}{120} \tau^5 \right) - \frac{1}{360} \tau^5 \left( \frac{1}{K} \right)^4$$

Parisi continuous ansatz proven right

# SK model in magnetic field

- Full RSB at AT line reduces to 1RSB ( $h > 0$ )
- Small expansion parameter  $\alpha = \beta^2 \langle (1 - t_0^2)^2 \rangle_\eta - 1$   
( $t_0 \equiv \tanh[\beta(h + \eta\sqrt{q})]$ )

$$\nu = \frac{2 \langle t_0^2 (1 - t_0^2)^2 \rangle_\eta}{\langle (1 - t_0^2)^3 \rangle_\eta}$$

$$\chi_1 = \frac{1}{2\beta^2\nu} \frac{\beta^2 \langle (1 - t_0^2)^2 \rangle_\eta - 1}{1 - 3\beta^2 \langle t_0^2 (1 - t_0^2)^2 \rangle_\eta} + O(\alpha^2)$$

$$\nu_l^K \doteq \nu_1 + (K+1-2l)\Delta\nu/K, \quad \Delta\chi_l^K \doteq \chi_1/K$$

$$\Delta\nu \doteq \frac{\beta^2 \chi_1 \left\langle (1 - t_0^2)^2 \left( 2(1 - 3t_0^2)^2 + 3(t_0^2 - 1)\nu(8t_0^2 + (t_0^2 - 1)\nu) \right) \right\rangle_\eta}{\langle (1 - t_0^2)^3 \rangle_\eta}$$

$$\Lambda_l^K \doteq - \frac{2\beta^2}{3K^2} \frac{\chi_1 \Delta\nu}{\nu + 2}$$

# Potts glass ( $p < 4$ ): discrete RSB

- Two 1RSB solutions for  $\nu_1 \doteq \frac{p-2}{2} + \frac{36-12p+p^2}{8(4-p)}\tau$

- Locally stable solution (near  $T_c$  and  $p > p^* \approx 2.82$ )

$$q^{(1)} \doteq 0, \quad \Delta\chi^{(1)} \doteq \frac{2}{4-p}\tau$$

Stability function:  $\Lambda_1^{(1)} \doteq \frac{\tau^2(p-1)}{6(4-p)^2} (7p^2 - 24p + 12)$

- Unstable solution ( $p > p^*$  unphysical)

$$q^{(2)} \doteq \frac{-12 + 24p - 7p^2}{3(4-p)^2(p-2)}\tau^2, \quad \Delta\chi^{(2)} \doteq \frac{2}{4-p}\tau$$

- $K$  RSB (from the unstable one)


$$q^K \doteq -\frac{1}{3K^2} \frac{12 - 24p + 7p^2}{(4-p)^2(p-2)}\tau^2, \quad \Delta\chi_l^K \doteq \frac{1}{K} \frac{2}{(4-p)}\tau,$$

$$\nu_l^K \doteq \frac{p-2}{2} + \frac{2}{4-p} \left[ 3 + \frac{3}{2}p - p^2 + \left( 3 - 6p + \frac{7}{4}p^2 \right) \frac{2l-1}{2K} \right] \tau$$

Copyright © 2015, All rights reserved. http://www.cmapx.jku.cz/~jancik


# Potts glass ( $p = 3$ ): coexistence

1RSB and FRSB coexist near  $T_c$



Stability and entropy of 1RSB solution ( $p = 3$ )  
Free-energy differences:

$$\beta(f_c - f_{1RSB}) \doteq \frac{(p-1)(p(7p-24)+12)^2 \tau^5}{720(4-p)^5}, \quad \beta(f_c - f_{RS}) \doteq \frac{(p-1)(p-2)^2 \tau^3}{3(4-p)(6-p)^2}$$



Free-energy difference as function of  $p$

# p-spin glass: 1RSB 1

- Discontinuous transition to the low-temperature phase for  $p > 2$
- Asymptotic solution  $p \rightarrow \infty$ : 1RSB

$$\begin{aligned}
 f_T^{(p \rightarrow \infty)}(q, \chi_1, \mu_1) &= -\frac{1}{4T} [1 - (q + \chi_1)(1 - \ln(q + \chi_1))] - \frac{1}{\mu_1} \ln [2 \cosh(\mu_1 h)] \\
 &\quad - \frac{\mu_1}{4} [\chi_1 - (q + \chi_1) \ln(q + \chi_1)] - \frac{\mu_1 q}{4} \left[ \ln q + p \left( 1 - \tanh^2(\mu_1 h) \right) \right]
 \end{aligned}$$

rescaled variable  $\mu_1 = \beta v_1$

- Low-temperature solution ( $p = \infty$ ) -- Random energy model

$$\begin{aligned}
 \chi_1 &= 1 - q, \quad q = \exp\{-p(1 - \tanh^2(\mu_1 h))\}, \\
 \mu_1 &= 2\sqrt{\ln [2 \cosh(\mu_1 h)] - h \tanh(\mu_1 h)}
 \end{aligned}$$

for  $\beta > 2\sqrt{\ln [2 \cosh(\beta h)] - h \tanh(\beta h)}$ ,

otherwise  $q + \chi_1 = 0$  and  $\mu_1 = \beta$



# Conclusions

## Ergodicity breaking without symmetry breaking

- 1 Frustration with disorder prevents existence of physical symmetry-breaking fields
- 2 **Real replicas** -- means to test thermodynamic homogeneity (ergodicity)
- 3 **Analytic continuation to non-integer replication index mandatory** -- ultrametric structure
- 4 Broken LRT of inter-replica interaction -- broken replica symmetry (ergodicity)
- 5 Hierarchical replications -- series of admissible solutions (equilibrium states)
- 6 Local and global stability conditions select the true equilibrium
- 7 **Continuous RSB -- marginally stable (available only via asymptotic expansions)**